



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

Let $\rho \equiv r/R$ so that the cubic becomes

$$\rho^3 - \rho^2 + 1/(3n) = 0. \quad (3)$$

Since the constant term of this equation is positive there is always one real negative root, and this is inadmissible. Therefore the remaining roots must be real in order that a positive root may exist. The derivative of the left-hand member of (3) is $\rho(3\rho - 2)$ which vanishes for $\rho = 0$ and $\rho = 2/3$. The expression for the derivative shows that there is a maximum for $\rho = 0$, and a minimum for $\rho = 2/3$; the value of the minimum is $1/3(1/n - 4/9)$ and hence if $1/n > 4/9$ there is only one real root, the negative root already mentioned.

Consequently the given value $1/n = 3/4$ conflicts with the preceding condition and makes the problem impossible.

Also solved by T. M. BLAKSLER, L. A. EASTBURN, R. A. JOHNSON, R. H. MARSHALL, J. Q. McNATT, ARTHUR PELLETIER, and A. V. RICHARDSON.

2855 [1920, 377]. Proposed by J. L. RILEY, Stephenville, Texas.

Show that the circle of curvature at any point of the ellipse cannot pass through the centre unless the eccentricity be greater than $1/\sqrt{2}$.

I. SOLUTION BY A. V. RICHARDSON, Bishop's College, Lennoxville, Quebec.

In the ellipse

$$\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1 \quad (a > b),$$

the normal at the point $(a \cos \theta, b \sin \theta)$ is

$$\frac{ax}{\cos \theta} - \frac{by}{\sin \theta} = a^2 - b^2. \quad (1)$$

To find the coördinates of the center of curvature, we solve (1) and its derivative with respect to θ ,

$$\frac{ax \sin \theta}{\cos^2 \theta} + \frac{by \cos \theta}{\sin^2 \theta} = 0,$$

giving

$$x = \frac{a^2 - b^2}{a} \cos^3 \theta; \quad y = -\frac{a^2 - b^2}{b} \sin^3 \theta. \quad (2)$$

The (radius of curvature)² is, of course, the (distance)² between this point and $(a \cos \theta, b \sin \theta)$ i.e.,

$$\frac{(a^2 \sin^2 \theta + b^2 \cos^2 \theta)^3}{a^2 b^2}.$$

If we put the radius of curvature equal to the distance of the center of curvature from the origin we shall get after some reductions

$$(a^2 - 2b^2) \cos^2 \theta = (2a^2 - b^2) \sin^2 \theta. \quad (3)$$

Hence, for real values of θ , we must have $a^2 > 2b^2$, i.e.,

$$a^2 > 2a^2(1 - e^2) \quad \text{or} \quad e > \frac{1}{\sqrt{2}}.$$

II. REMARKS BY OTTO DUNKEL, Washington University.

If we determine the condition that the perpendicular bisector of the segment from the origin to the point $(a \cos \theta, b \sin \theta)$ shall pass through the center of curvature we shall get equation (3) more easily.

Or we may express this by saying that the center of curvature projects into the middle point of the segment. This condition takes the form $2(x_1x_c + y_1y_c) - (x_1^2 + y_1^2) = 0$, which reduces to the same equation (3).

We find also that e may equal $1/\sqrt{2}$. In the ellipse for which $e = 1/\sqrt{2}$ the radius of curvature at the extremity of the major axis is equal to $a/2$.

Also solved by A. M. HARDING, WILLIAM HOOVER, and H. S. UHLER.